

Comparing Effectiveness of Family Planning Methods

More effective


Less than 1 pregnancy per 100 women in 1 year



Implants




IUD



Female sterilization




Vasectomy



Injectables




LAM




Pills



Patch



Vaginal ring



Male condoms



Diaphragm



Female condoms



Fertility awareness methods



Withdrawal



Spermicides

Less effective

About 30 pregnancies per 100 women in 1 year

How to make your method more effective

Implants, IUD, female sterilization: After procedure, little or nothing to do or remember

Vasectomy: Use another method for first 3 months

Injectables: Get repeat injections on time

Lactational amenorrhea method, LAM (for 6 months): Breastfeed often, day and night

Pills: Take a pill each day

Patch, ring: Keep in place, change on time

Condoms, diaphragm: Use correctly every time you have sex

Fertility awareness methods: Abstain or use condoms on fertile days. Newest methods (Standard Days Method and TwoDay Method) may be easier to use.

Withdrawal, spermicides: Use correctly every time you have sex



Sources:

Steiner MJ, Trussell J, Mehta N, Condon S, Subramaniam S, Bourne D. Communicating contraceptive effectiveness: a randomized controlled trial to inform a World Health Organization family planning handbook. *Am J Obstet Gynecol* 2006;195:85–91.

World Health Organization Department of Reproductive Health and Research (WHO/RHR), Johns Hopkins Bloomberg School of Public Health (JHSPH)/Center for Communication Programs (CCP). *Family Planning: A Global Handbook for Providers*. Baltimore, MD and Geneva: CCP and WHO, 2007.

Trussell J. Choosing a contraceptive: efficacy, safety, and personal considerations. In: Hatcher RA, Trussell J, Stewart F, Nelson AL, Cates W Jr., Guest F, Kowal D, eds. *Contraceptive Technology, Nineteenth Revised Edition*. New York: Ardent Media, Inc., in press.